

Relationship of alcohol intake and thiamine deficiency in heart failure

ASA van der Werff¹, A Klooster^{2*}

Abstract

Introduction

The purpose of this paper is to summarize the relationship of alcohol intake with thiamine deficiency and with heart failure and the manner in which they interact with each other.

Discussion

Moderate alcohol consumption has shown to decrease cardiovascular mortality. However, heavy alcohol consumption is associated with alcoholic cardiomyopathy. Excessive alcohol intake is associated with thiamine deficiency. Moreover, heart failure is also related to thiamine deficiency, and thiamine supplementation has shown to increase left ventricular ejection fraction.

Conclusion

Thiamine deficiency should be further studied in heart failure, and more research is needed to show the effect of thiamine supplementation in heart failure.

Introduction

Excessive alcohol use is the third leading lifestyle-related cause of death for people in the United States each year, placed behind tobacco and improper diet/lack of physical activity, which are the first and second lifestyle-related causes of death, respectively¹.

Alcohol in alcoholic drinks is mainly ethanol. The ethanol molecule can, because of its structure and small size, diffuse readily across

cell membranes. Because there is no identifiable receptor, the mode of action is non-specific. The effects of ethanol are complex and contain one or several of the following mechanisms: (1) changes in membrane fluidity, (2) changes in the relative concentration or organization of specific membrane lipids, (3) alterations in specific protein domains (that in turn alter membrane protein function), (4) stimulation or attenuation of membrane signal transduction cascades and related intra-cellular enzyme systems (e.g. protein kinase A and C), (5) changes in channel function, (6) changes in receptor number and subunit expression and finally (7) modulation of neurotransmitters and neuromodulator systems².

However, besides these rather detrimental effects, it has extensively been shown that in apparently healthy subjects, moderate alcohol consumption (1–2 drinks/day) decreases cardiovascular and all-cause mortality³. Also, the risk of coronary heart disease, ischaemic strokes and amputations due to peripheral vascular disease is decreased with moderate alcohol consumption⁴. Mechanisms supporting this positive effect of moderate alcohol consumption include beneficial regulation of lipids and fibrinolysis, decreased platelet aggregation and coagulation factors and beneficial effects on endothelial function, inflammation and insulin resistance³.

The relationship between alcohol and total mortality appears to be J-shaped⁵. Consequently, it has been a matter of great debate on what amount and duration of alcohol abuse is required to produce detrimental effects. It appears that

most alcoholic patients with detectable changes in cardiac structure and function report consuming >90 g alcohol per day for ≥5 years. Therefore, heavy drinking refers to more than six standard alcoholic drinks per day for more than 5 years.

Excessive alcohol intake is also related to thiamine deficiency. Thiamine deficiency, especially in relation to alcohol abuse, leads to Wernicke encephalopathy and Korsakoff syndrome⁶. Thiamine deficiency can also result in wet beriberi. Clinical features of wet beriberi will respond within hours to a bolus of thiamine⁷. There is increasing evidence that thiamine deficiency also plays a role in heart failure.

The purpose of this paper is to summarize the relationship of alcohol intake with thiamine deficiency and with heart failure and the way they interact with each other.

Discussion

Alcohol and heart failure

Lifestyle modification including moderation of alcohol intake is important in cardiovascular disease. Among heart failure patients, one in five patients drink two or more alcoholic beverages everyday⁸. This puts them at risk of alcohol dependence.

The acute effects of alcohol consumption are mainly obtained from isolated and whole animal heart preparations and young healthy human persons². Results of these studies are not clear, and the main limitation is that the sample sizes range from 4 to 12.

Greenberg et al.⁹ specifically researched the acute effects of alcohol consumption on haemodynamic and echocardiographic parameters in

* Corresponding author
Email: a.klooster@umcg.nl

¹ Department of Cardiology, Medisch Spectrum Twente, Enschede, The Netherlands.

² Department of Pathology and Medical Biology, University Medical Centre Groningen, Groningen, The Netherlands.

eight heart failure patients. Alcohol ingestion resulted in rather favourable haemodynamic effects, which include a significant reduction in mean arterial pressure, pulmonary capillary wedge pressure and systemic vascular resistance.

There is limited data on the long-term effects of moderate alcohol intake in heart failure patients. Cooper et al.¹⁰ specifically investigated the effects of moderate drinking in heart failure patients. It was shown that moderate drinking is associated with a significant reduction in relative risk for all-cause mortality and fatal myocardial infarction in the group with ischaemic left ventricular dysfunction. However, in the non-ischaemic left ventricular dysfunction group, moderate drinking had no effect on any of the outcome parameters.

In a meta-analysis on alcohol consumption and mortality in patients with cardiovascular disease (coronary heart disease, acute myocardial infarction and stroke), it was shown that there was a J-shaped relationship between alcohol consumption and all-cause and cardiovascular mortality³. It seems that the effect of moderate drinking is similar in patients with cardiovascular disease as in healthy subjects.

As moderate alcohol consumption appears to be safe and may also have beneficial effects, one could argue that moderate drinking should be recommended. However, there is currently insufficient evidence to recommend the use of alcohol in heart failure patients that do not currently consume drink alcohol. One reason to be conservative is that there is considerable individual variation in the effect of alcohol consumption. There is a three- to four-fold variation in the rate of alcohol metabolism between individuals, and there is a two- to three-fold variation in the pharmacodynamics of alcohol due to individual differences¹¹.

Long-term heavy alcohol consumption is known to lead to

non-ischaemic dilated cardiomyopathy, which is referred to as alcoholic cardiomyopathy if heavy drinking is reported. Alcoholic cardiomyopathy is responsible for 21%–36% of all cases of non-ischaemic dilated cardiomyopathy in Western Society. The prevalence of alcoholic cardiomyopathy among alcoholics is variable and ranges from 23% to 40%¹¹. It occurs more in men than women, although women are more sensitive to the toxic dose of alcohol. Limited data are available on the amount and duration of consumption required to produce symptomatic alcoholic cardiomyopathy. However, in general, heavy drinking (>90 g alcohol per day, which equals six standard alcoholic drinks per day) for more than 5 years puts patients at risk for the development of asymptomatic alcoholic cardiomyopathy².

Clinical findings in patients with alcoholic cardiomyopathy are identical to idiopathic cardiomyopathy, with dilatation and impaired contraction of the left or both ventricles, increased left ventricular end-diastolic volume, increased left ventricular mass and a left ventricular ejection fraction below 45%¹². Therefore, the diagnosis of alcoholic cardiomyopathy is based on the coincidence of heavy alcohol consumption and a global myocardial dysfunction with no other underlying myocardial disease. However, alcoholic cardiomyopathy is not necessarily associated with deterioration after ceasing alcohol consumption while the idiopathic form is¹². If patients with alcoholic cardiomyopathy do not abstain from alcohol, they have a poorer prognosis and significantly higher mortality rates compared with those who do abstain. The 4-year mortality can be as high as 50%^{2,13}.

Alcohol and thiamine

Alcohol consumption leads to thiamine deficiency by decreasing the transport of thiamine across the intestinal mucosa, impairing capacity

of the liver to store thiamine and impairing the phosphorylation of thiamine to the more biological active thiamine diphosphate¹⁴.

The active form of thiamine, thiamine diphosphate, is the co-enzyme of three major enzymes: transketolase (TK), pyruvate dehydrogenase complex (PDH) and α -ketoglutarate dehydrogenase (α -KHD). These enzymes are present in the mitochondria and are essential in citric acid cycle and for the generation of ATP. Alcohol especially decreases the amount of thiamine diphosphate. *In vitro* and *in vivo* experiments have shown that thiamine pyrophosphokinase is inhibited by ethanol¹⁵. Furthermore, chronic ethanol intake leads to increased thiamine diphosphatase¹⁶. As shown in Figure 1, this double action, due to chronic exposure to ethanol, further decreases the amount of thiamine diphosphate.

Moreover, the metabolism of alcohol raises the demand for thiamine. In this way, alcohol-dependent people require more thiamine than non-alcoholic people. Other contributing factors to thiamine deficiency in alcoholics are self-neglect and low content of vitamins and minerals of alcoholic beverages.

Thiamine deficiency in heart failure patients

Thiamine deficiency is commonly linked to alcoholism, but can be seen in other patients as well. Special attention is paid in this respect to people with a staple diet of white bread or polished rice^{17,18}. But, even more recently, thiamine deficiency has been described in several populations that rely entirely on the provision of nutrition by outside sources^{19,20}. Also, parenteral nutrition can be thiamine-deficient, which can cause an outbreak of thiamine deficiency in infants²¹.

Thiamine deficiency can lead to wet beriberi. The infantile form is well described in infants who were fed parenteral nutrition that




Figure 1: Thiamine phosphorylation and dephosphorylation. Influence of alcohol is represented with grey arrows, resulting in decreased thiamine diphosphate (TDP).

appeared to be thiamine-deficient²¹. Symptoms in adults can be divided into a common high output state and an uncommon low output state. Symptoms of wet beriberi in infants and adults are summarized in Table 1.

Among patients with heart failure, thiamine deficiency is highly prevalent. Potential contributing factors of thiamine deficiency in patients with heart failure are loop diuretic use, malnutrition, advanced age, heart failure severity and frequent hospitalization. Incidence of thiamine deficiency in heart failure patients has been reported to be 33%, whereas thiamine deficiency was present in 12% of the control subjects²². In this study by Hanninen et al., thiamine deficiency was related to urine thiamine loss, absence of intake of thiamine-containing supplements and preserved renal function.

Furosemide treatment has shown to exacerbate the thiamine deficiency, probably by preventing the re-absorption of thiamine and thereby increasing the urinary excretion of thiamine²³. Digoxin has been shown to have a similar effect, and furosemide and digoxin have an additive effect if used simultaneously²⁴.

By Zenuk et al.²⁵ it was shown that with furosemide therapy, at doses higher than 80 mg/day, prevalence of thiamine deficiency was 98%. However, thiamine deficiency may also occur with long-term use of smaller dosages of diuretics.

In vitro experiments showed that thiamine supplementation has a cytoprotective effect on cardiomyocytes during hypoxic insult²⁶. Hypoxia is an important cause of cell death in myocardial infarction, cardiomyopathy and chronic heart failure. It has also been shown that the cardiac contractility is reduced by thiamine deficiency²⁷.

Until now, four studies examined the effect of thiamine supplementation on cardiac function in heart failure patients²⁸⁻³¹. Three of these studies, of which two were randomized, found an improvement of left ventricular function after thiamine supplementation²⁸⁻³⁰. One study did not show a significant effect of thiamine on the clinical course of cardiomyopathy, but cardiac function was not assessed by echocardiography³¹. The most recent study was a randomized, double-blind, placebo-controlled, crossover pilot study in nine patients²⁸. After 28 days of treatment with 300 mg thiamine a day, the left ventricular ejection fraction was significantly increased.

Table 1 Clinical co-features of thiamine deficiency related to different forms of beriberi

Infantile beriberi	
Cardiomyopathy	
Aphonia	
Absent deep-tendon reflexes	
Vomiting	
Diarrhoea	
Weight loss	
Restlessness	
Nystagmus	
Ophthalmoplegia	
Respiratory symptoms	
Wet beriberi	
<i>Common high output state</i>	<i>Uncommon low output state</i>
Heart failure	Severe hypotension
Orthopnea	Lactic acidosis
Pulmonary oedema	Absence of oedema
Peripheral oedema	

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: van der Werff ASA, Klooster A. Relationship of alcohol intake and thiamine deficiency in heart failure. OA Alcohol 2013 Mar 01;1(1):4.

In a small observational study, it was shown that thiamine levels decreased during coronary artery bypass graft. The investigators propose that increased metabolic demand will increase the usage of thiamine in “stress situations” and will lead to increased depletion.

Interaction between alcohol, thiamine deficiency and heart failure

In Figure 2, the schematic relationship of alcohol, thiamine deficiency and heart failure is depicted. Whereas moderate alcohol intake seems to decrease cardiovascular and all-cause mortality, prolonged alcohol abuse results in alcoholic cardiomyopathy. Furthermore, alcohol is directly toxic to the cardiomyocyte. Therefore, alcohol abuse will lead to heart failure. Alcohol abuse also results in thiamine deficiency by improper intake of thiamine through self-neglect, decreased thiamine diphosphate, as a direct effect of alcohol and raised demand for thiamine.

Also, heart failure and thiamine deficiency interplay together. Heart failure patients are prone to thiamine deficiency by their general health factors (such as malnutrition and advanced age), and also treatment with furosemide and digoxin has

shown to increase thiamine deficiency. Thiamine deficiency was shown to decrease the left ventricular ejection fraction, thereby requiring more intensive treatment and probably worsening of the thiamine deficiency.

Conclusion

Heart failure and thiamine deficiency are closely linked not only because of the effect of alcohol on both of them but also because of their interplay together. Therefore, thiamine deficiency should be further studied in heart failure, and more research is needed to show the effect of thiamine supplementation in heart failure.

References

1. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA*. 2004 Mar;291(10):1238–45.
2. Piano MR. Alcohol and heart failure. *J Cardiac Fail*. 2002 Aug;8(4):239–46.
3. Costanzo S, Di Castelnuovo A, Donati MB, Iacoviello L, de Gaetano G. Alcohol consumption and mortality in patients with cardiovascular disease: a meta-analysis. *J Am Coll Cardiol*. 2010 Mar;55(13):1339–47.
4. Beulens JW, Algra A, Soedamah-Muthu SS, Visseren FL, Grobbee DE, et al. Alcohol consumption and risk of recurrent cardiovascular events and mortality in patients with clinically manifest vascular disease and diabetes mellitus: the second manifestations of ARterial (SMART) disease study. *Atherosclerosis*. 2010 Sep; 212(1):281–6.
5. Di Castelnuovo A, Costanzo S, Bagnardi V, Donati MB, Iacoviello L, et al. Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34 prospective studies. *Arch Intern Med*. 2006 Dec;166(22):2437–45.
6. Harper C. Thiamine (vitamin B1) deficiency and associated brain damage is still common throughout the world and prevention is simple and safe! *Eur J Neurol*. 2006 Oct;13(10):1078–82.
7. Naidoo DP, Gathiram V, Sadhabiriss A, Hassen F. Clinical diagnosis of cardiac beriberi. *South Afr Med J*. 1990 Feb; 77(3):125–7.
8. Marti CN, Georgiopoulou VV, Giamouzis G, Cole RT, Deka A, et al. Patient-reported selective adherence to heart failure self-care recommendations: a prospective cohort study: the Atlanta Cardiomyopathy Consortium. *Congest Heart Fail*. 2013 Jan-Feb;19(1):16–24.
9. Greenberg BH, Schutz R, Grunkemeier GL, Griswold H. Acute effects of alcohol in patients with congestive heart failure. *Ann Intern Med*. 1982 Aug;97(2):171–5.
10. Cooper HA, Exner DV, Domanski MJ. Light-to-moderate alcohol consumption and prognosis in patients with left ventricular systolic dysfunction. *J Am Coll Cardiol*. 2000 Jun;35(7):1753–9.
11. Iacovoni A, De Maria R, Gavazzi A. Alcoholic cardiomyopathy. *J Cardiovasc Med*. 2010 Dec;11(12):884–92.
12. Schoppet M, Maisch B. Alcohol and the heart. *Herz*. 2001 Aug;26(5):345–52.
13. George A, Figueredo VM. Alcoholic cardiomyopathy: a review. *J Cardiac Fail*. 2011 Oct;17(10):844–9.
14. Thomson AD. Mechanisms of vitamin deficiency in chronic alcohol misusers and the development of the wernicke-korsakoff syndrome. *Alcohol Alcohol*. 2000 May-Jun;Suppl 35(1):2–7.
15. Rindi G, Imarisio L, Patrini C. Effects of acute and chronic ethanol administration on regional thiamin pyrophosphokinase activity of the rat brain. *Biochem Pharmacol*. 1986 Nov; 35(22):3903–8.
16. Laforenza U, Patrini C, Gastaldi G, Rindi G. Effects of acute and chronic ethanol administration on thiamine metabolizing enzymes in some brain areas and in other organs of the rat. *Alcohol Alcohol*. 1990;25(6):591–603.




Figure 2: Schematic relationship of alcohol, thiamine deficiency and heart failure.

Licensee OA Publishing London 2013. Creative Commons Attribution License (CC-BY)

FOR CITATION PURPOSES: van der Werff ASA, Klooster A. Relationship of alcohol intake and thiamine deficiency in heart failure. *OA Alcohol* 2013 Mar 01;1(1):4.

17. Doung-ngern P, Kesornsukhon S, Kanyanaphotporn J, Wanadurongwan S, Songchitsomboon S. Beriberi outbreak among commercial fishermen, Thailand 2005. *Southeast Asian J Trop Med Public Health*. 2007 Jan;38(1):130–5.
18. Kamien M, Nobile S, Cameron P, Rosevear P. Vitamin and nutritional status of a part aboriginal community. *Aus N Z J Med*. 1974 Apr;4(2):126–37.
19. Ahoua L, Etienne W, Fermon F, Godain G, Brown V, et al. Outbreak of beriberi in a prison in cote d'ivoire. *Food Nutr Bull*. 2007 Sep;28(3):283–90.
20. Watson JT, El Bushra H, Lebo EJ, Bwire G, Kiyengo J, et al. Outbreak of beriberi among african union troops in mogadishu, Somalia. *PloS One*. 2011; 6(12):e28345.
21. Fattal-Valevski A, Kesler A, Sela BA, Nitzan-Kaluski D, Rotstein M, et al. Outbreak of life-threatening thiamine deficiency in infants in israel caused by a defective soy-based formula. *Pediatrics*. 2005 Feb;115(2):e233–8.
22. Hanninen SA, Darling PB, Sole MJ, Barr A, Keith ME. The prevalence of thiamin deficiency in hospitalized patients with congestive heart failure. *J Am Coll Cardiol*. 2006 Jan;47(2):354–61.
23. Yui Y, Itokawa Y, Kawai C. Furosemide-induced thiamine deficiency. *Cardiovasc Res*. 1980 Sep;14(9):537–40.
24. Zangen A, Botzer D, Zangen R, Shainberg A. Furosemide and digoxin inhibit thiamine uptake in cardiac cells. *Eur J Pharmacol*. 1998 Nov;361(1):151–5.
25. Zenuk C, Healey J, Donnelly J, Vaillancourt R, Almalki Y, et al. Thiamine deficiency in congestive heart failure patients receiving long term furosemide therapy. *Can J Clin Pharmacol*. 2003 Winter; 10(4):184–8.
26. Shin BH, Choi SH, Cho EY, Shin MJ, Hwang KC, et al. Thiamine attenuates hypoxia-induced cell death in cultured neonatal rat cardiomyocytes. *Mol Cells* 2004 Oct;18(2):133–40.
27. Cappelli V, Bottinelli R, Polla B, Reggiani C. Altered contractile properties of rat cardiac muscle during experimental thiamine deficiency and food deprivation. *J Mol Cell Cardiol*. 1990 Oct; 22(10):1095–106.
28. Schoenenberger AW, Schoenenberger-Berzins R, der Maur CA, Suter PM, Vergopoulos A, et al. Thiamine supplementation in symptomatic chronic heart failure: A randomized, double-blind, placebo-controlled, cross-over pilot study. *Clin Res Cardiol*. 2012 Mar;101(3):159–64.
29. Seligmann H, Halkin H, Rauchfleisch S, Kaufmann N, Motro M, et al. Thiamine deficiency in patients with congestive heart failure receiving long-term furosemide therapy: a pilot study. *Am J Med*. 1991 Aug;91(2):151–5.
30. Shimon I, Almog S, Vered Z, Seligmann H, Shefi M, et al. Improved left ventricular function after thiamine supplementation in patients with congestive heart failure receiving long-term furosemide therapy. *Am J Med*. 1995 May; 98(5):485–90.
31. Pfitzenmeyer P, Guillard JC, d'Athis P, Petit-Marnier C, Gaudet M. Thiamine status of elderly patients with cardiac failure including the effects of supplementation. *Int J Vit Nutr Res*. 1994; 64(2):113–8.