

Rotator cuff tears, evaluation and treatment: a critical review

B Haviv^{1,2*}

Abstract

Introduction

Rotator cuff tears are a common cause for shoulder pain in the older population. The aetiology and pathophysiology are not fully understood. Theories for tear evolvment are divided into intrinsic (e.g. recurrent microtrauma) and extrinsic (e.g. subacromial impingement). The subacromial bursa is probably the source of pain in symptomatic patients with rotator cuff tears. It is uncertain whether the tear itself can produce pain because of the high prevalence of asymptomatic tears. Typically, there is a gradual increase in shoulder pain and weakness; however, it can present acutely due to an injury. Difficulties in overhead activities and night pain are common. Imaging such as ultrasound and magnetic resonance are required for accurate evaluation of RCTs in patients with ongoing pain and limitations. Primary treatment includes activity modifications, pain relief and physical therapy. Surgery is advised for acute tears in active patients or chronic symptomatic tears in patients that fail to improve. This article discusses the evaluation and treatment of rotator cuff tears.

Conclusion

Symptoms of pain and weakness about the shoulder imply on rotator cuff tear. There are typical

impingement signs in physical evaluation together with limited range of motion and weakness. Plain radiography is required to rule out other pathologies and observe acromion morphology, while ultrasound and magnetic resonance imaging are used to define the tear and tendon quality before surgical intervention. Most repairs are currently performed in an all-arthroscopic minimally invasive technique with easier rehabilitation and less pain compared to the traditional open surgery.

Introduction

Shoulder pain is a very common complaint. Studies report on about one-third of the population that suffer from shoulder symptoms during their lifetime. One of the main causes for shoulder pain in the older population is rotator cuff tears (RCTs). The prevalence of RCTs in the general population is 20%¹. Without appropriate evaluation and treatment, the pain may persist for a long period of time. Since the early 1990s, there was a significant progress in arthroscopic surgery techniques for symptomatic tears of the rotator cuff. Currently, selected patients are treated with these techniques with high success and low morbidity rates. The aim of this review was to discuss the methods for evaluation and the treatment techniques for RCTs.

Anatomy

There are two synovial spaces in the shoulder—the glenohumeral and subacromial spaces. Between those spaces is the rotator cuff complex of four tendons together with the underlined joint capsule². These are the supraspinatus, infraspinatus and

teres minor, which originate from the posterior scapula and insert into the greater tuberosity of the proximal humerus, and the subscapularis that originate from the anterior scapula and insert into the lesser tuberosity. The RC moves and stabilises the humeral head in the centre of the glenoid by the principle of coupling forces. The subacromial bursa is localised in between the RC below and the acromion with the coracoacromial ligament (CAL) above. The normal separation between the glenohumeral and subacromial spaces is violated when there is a full-thickness tear of the RC. The most common tear of the RC involves the supraspinatus tendon. This tendon has a unique structure of several parallel independent fibre units that allows for wide-ranging mobility³. During movement, some units are elongated, while others are shortened. The sheering forces that are produced can cause pathology.

Tear types

Accurate description of RCTs is essential for clinical and academic purposes. Measurements can be done by imaging or at surgery. It is usually defined by the tendon involved, its thickness (partial or complete) and size (anterior to posterior). Crescent shape tear is the most common full-thickness configuration.

Pathophysiology

Codman⁴ described a hypovascularised zone adjacent to the supraspinatus insertion site. Although the pathological process is not fully understood, it is degenerative and deteriorates with age. Imaging studies have shown that the prevalence of asymptomatic RCTs is 30% and 65%

* Corresponding author
Email: barak_haviv69@hotmail.com

¹ Arthroscopy and Sports Injuries Unit, Hasharon Hospital, Rabin Medical Center, Petach-Tikva, Israel

² Orthopedic Department, Sackler Faculty of Medicine, Tel-Aviv University P.O. Box 39040, Tel Aviv 6997801, Israel

in people older than 60 and 70 years, respectively⁵.

Traditionally, the pathophysiological theories are divided into intrinsic and extrinsic.

Intrinsic tendinopathy

The pathological process involves the tendon tissue. The main model is degenerative with age-related deterioration into partial- and then full-thickness tendon rupture. Because of recurrent micro-trauma, inflammatory mediators alter the local environment and oxidative stress causes tenocyte loss with further degeneration. Both histological examinations of tendon tissue and overload animal models have demonstrated changes in blood supply and also in concentrations of cells, collagen and proteoglycans. In addition, studies have shown loss of fibrous organisation, tissue weakening and high level of inflammatory markers⁶⁻⁹. Shoulder abduction causes a relatively high mechanical stresses in the articular side compared to bursal side fibres of the supraspinatus tendon¹⁰. These forces produce cartilage tissue near the insertion of the supraspinatus tendon into the greater tuberosity as was demonstrated in other parts of the body¹¹. This cartilaginous tissue has weaker load resistance.

Burkhart et al.¹² have described the 'rotator cuff cable', a thick crescent-like structure adjunct to humeral insertion site of the RC. The cable is perpendicular to the tendon fibres where the medial fibres between it and the humerus are thinner and localised in the hypovascular critical zone described by Codman⁴. Although the cable has biomechanical advantages derived from the suspension bridge mechanical principle, these medial fibres tend to wear and tear.

The vascular supply to the supraspinatus tendon has been studied for many years. The common theory suggests a critical hypovascular zone at 10–15mm from its insertion site

into the humerus. However, several studies have contradicted this theory, and thus the relationship between vascular supply and degeneration is not entirely clear¹³. Nevertheless, the tear margins are often avascular with limited healing potential.

Extrinsic tendinopathy

The pathological process is external to the tendon tissue. Neer has suggested the coracoacromial arch as the primary offender¹⁴. The recurrent attrition of the tendons against the inferior part of an aberrant acromion causes tissue damage. According to Bigliani et al.¹⁵, there are three morphological types of acromion. Of these, curved or hooked acromion morphologies are found in most patients with RCTs. Based on the above assumptions, one of the most popular surgical techniques of the shoulder is acromioplasty (resurfacing the lower edge of the acromion), although its correlation with clinical results is currently debatable¹⁶. Pain receptors have been found at the CAL, and some authors believe that it may cause external impingement¹⁷, while others claim that the pathological changes within the CAL are secondary to chronic mechanical tension¹⁸ and do not cause RCTs. Surgical detachment of the CAL may cause superior migration of the humeral head, and therefore it is important to try and preserve it.

Theoretically, subacromial impingement should trigger a partial tear at the subacromial tendon side; however, there are many reports on partial supraspinatus tear at its articular side¹⁹. This has led to the internal (or superior posterior) impingement theory in which the RC is compressed between the posterior glenoid margins and the greater tuberosity.

Several demographic factors have been implied to be correlated with RCTs. For example, it would be logical to assume higher tear rates in the dominant shoulder; still, about

one-third of the patients with symptomatic tears have asymptomatic contralateral full-thickness tears, and most patients with symptomatic tears are not manual labourers²⁰. Another example is the negative effect of smoking on tendon healing, which was evident after surgical repair and in animal studies²¹.

The source of pain

The source of pain in rotator cuff abnormalities is still unclear. Ruptured RC does not cause pain directly since there are many asymptomatic full-thickness tears. It is generally believed that the subacromial bursa is a major source of pain and discomfort as it undergoes some friction during shoulder movement and has sensory nerve endings²². In addition, the pain level was found to be correlated with the subacromial bursitis. The bursa is innervated anteriorly by the suprascapular nerve and posteriorly by the lateral pectoral nerve²³. It contains nociceptors and proprioceptors. These receptors and the presence of mechanoreceptors under the CAL imply a reflex system that coordinates the rotator cuff maintenance of humeral head position.

High levels of pain-associated proteins (e.g. substance P), inflammatory cytokines (e.g. IL1) and collagen-catabolic proteins were found in patients with shoulder impingement syndrome²⁴.

Diagnosis

The RC may rupture acutely due to trauma, but often the clinical presentation is gradual with progressive pain and weakness around the shoulder. Pain increases with shoulder elevation, internal rotation and at night. Shoulder strength depends on the tear size and the overall function of other intact tendons. Physical findings that are suggestive of tear are muscle atrophy, subacromial tenderness and crepitation. Strength and motion may be limited depending on the muscle involved.

Massive tears may cause shoulder instability, and thus when trying to elevate the arm, the humeral head subluxates anteriorly. In this position, the deltoid muscle cannot efficiently abduct the arm. This clinical presentation is called pseudo-paralysis. Imaging techniques are required if an acute tear is suspected or whenever the pain does not resolve. Plain radiographs are recommended as the primary modality to rule out other pathologies such as arthritis and to observe acromial morphology with its distance from the humeral head (decreased in massive tears); however, ultrasound (US) and magnetic resonance imaging (MRI) are used to define RCTs. Studies that have compared imaging modalities to arthroscopic findings have found similar accuracy of MRI and US in detecting RCTs with sensitivity of 97% and specificity of 67%²⁵.

Discussion

Treatment

Despite numerous publications regarding the treatment of RCTs, the number of high-level, evidence-based studies is scarce²⁶. Many studies are limited because of variations in control groups, outcome measures, short follow-ups and the ability to generalise outcomes of large referral medical centres to peripheral ones. Therefore, treatment is based on clinical experience, understanding the anatomy and management of other tendon ruptures in the body such as those of the hand.

Acute full-thickness tears

This should be repaired in up to 6 weeks from the injury, especially in relatively young and active patients; otherwise it might retract and atrophy with irreversible changes. If not repaired early, these tears may be hard to fix with bad outcome.

Partial tear

Many patients improve without surgery. Patients should be treated with exercises and stretches in order to

gain range of motion. Unresolved pain may benefit from arthroscopic debridement.

Chronic full-thickness tears

Many patients improve without surgery. Conservative treatment includes modifying activities, exercises, physical therapy, analgesics and non-steroidal anti-inflammatory medications. Subacromial steroid injections temporarily relieve pain. Recurrent injections should be avoided as they may impair tendon tissue, unless surgery is not considered and prior injections have helped.

Persistent pain and dysfunction may improve with surgical repair of the torn tendon. Factors that are correlated with good outcome after repair are age under 60 years, traumatic tear, short duration of symptoms (less than 2 months), no smoking, good general health, few injections (less than 4), primary repair, stable shoulder with good range of motion, no muscle atrophy, single tendon involvement (supraspinatus) and experienced surgeon²⁷.

The probability for recurrent tears after repairs increases with larger tears and atrophied tendons. The rehabilitation after surgery depends on the tear size and the quality of the repair. In many cases, active range of motion is forbidden for 6 or more weeks while the overall rehabilitation time may last longer than 6 months.

Surgical repair

Rotator cuff tendon repair is one of the most common procedures performed in the shoulder²⁸. During the past two decades, the traditional open approach has shifted to mini open and eventually to an all-arthroscopic minimally invasive technique²⁹. Open repairs have had good results but required partial deltoid muscle detachment. All arthroscopic techniques allow the treatment of concurrent intra-articular pathologies. It requires small incisions, less soft tissue

damage, no deltoid impairment and shorter postoperational pain and rehabilitation. Arthroscopic repairs have good long-term results but also have a steep learning curve.

Many studies have showed better results with non-recurrent (healed) tears after repairs. Therefore, there is much interest and recruitment of resources in order to achieve a successful repair by modifying important factors such as smoking cessation and enriching the biological environment at the tear site; however, the most dominant factor according to the literature is an anatomical stable fixation of the RCT³⁰.

Currently, an all-arthroscopic technique is utilised to perform anatomical anchoring of the rotator cuff to the proximal humerus. Common configurations of fixations are single or double row. Double-row fixations were found to have biomechanical advantages over single-row fixations; yet, there were no differences in clinical outcomes³⁰.

Conclusion

Symptoms of pain and weakness about the shoulder imply on RCT. There are typical impingement signs in physical evaluation together with limited range of motion and weakness. Plain radiography is required to rule out other pathologies and observe acromion morphology while US and MRI are used to define the tear and tendon quality before surgical intervention. The primary treatments involve activity modification, pain relief and exercises for a few months before considering surgical repair followed by prolonged rehabilitation. Acute repairs are suggested in traumatic tears in relatively young and active patients. At present, an all-arthroscopic technique is usually used to repair RCTs with good long-term results.

References

1. Yamamoto A, Takagishi K, Osawa T, Yanagawa T, Nakajima D, Shitara H, et al. Prevalence and risk factors of a rotator

- cuff tear in the general population. *J Shoulder Elbow Surg.* 2010 Jan;19(1):116–20.
2. Clark JM, Harryman DT 2nd. Tendons, ligaments, and capsule of the rotator cuff. Gross and microscopic anatomy. *J Bone Joint Surg Am.* 1992 Jun;74(5):713–25.
 3. Fallon J, Blevins FT, Vogel K, Trotter J. Functional morphology of the supraspinatus tendon. *J Orthop Res.* 2002 Sep;20(5):920–6.
 4. Codman E. The shoulder: Rupture of the supraspinatus tendon and other lesions in or about the subacromial bursa. Boston: Thomas Todd Company; 1934.
 5. Yamaguchi K, Ditsios K, Middleton WD, Hildebolt CF, Galatz LM, Teefey SA. The demographic and morphological features of rotator cuff disease. A comparison of asymptomatic and symptomatic shoulders. *J Bone Joint Surg Am.* 2006 Aug;88(8):1699–704.
 6. Hashimoto T, Nobuhara K, Hamada T. Pathologic evidence of degeneration as a primary cause of rotator cuff tear. *Clin Orthop Relat Res.* 2003 Oct;(415):111–20.
 7. Riley GP, Harrall RL, Constant CR, Chard MD, Cawston TE, Hazleman BL. Tendon degeneration and chronic shoulder pain: changes in the collagen composition of the human rotator cuff tendons in rotator cuff tendinitis. *Ann Rheum Dis.* 1994 Jun;53(6):359–66.
 8. Soslowsky LJ, Thomopoulos S, Tun S, Flanagan CL, Keefer CC, Mastaw J, et al. Overuse activity injures the supraspinatus tendon in an animal model: a histologic and biomechanical study. *J Shoulder Elbow Surg.* 2000 Mar–Apr;9(2):79–84.
 9. Perry SM, McIlhenny SE, Hoffman MC, Soslowsky LJ. Inflammatory and angiogenic mRNA levels are altered in a supraspinatus tendon overuse animal model. *J Shoulder Elbow Surg.* 2005 Jan–Feb;14(1 Suppl S):79S–83S.
 10. Reilly P, Amis AA, Wallace AL, Emery RJ. Mechanical factors in the initiation and propagation of tears of the rotator cuff. Quantification of strains of the supraspinatus tendon in vitro. *J Bone Joint Surg Br.* 2003 May;85(4):594–9.
 11. Lyman J, Weinhold PS, Almekinders LC. Strain behavior of the distal Achilles tendon: implications for insertional achilles tendinopathy. *Am J Sports Med.* 2004 Mar;32(2):457–61.
 12. Burkhart SS, Esch JC, Jolson RS. The rotator crescent and rotator cable: an anatomic description of the shoulder's "suspension bridge". *Arthroscopy.* 1993;9(6):611–6.
 13. Hegedus EJ, Cook C, Brennan M, Wyland D, Garrison JC, Driesner D. Vascularity and tendon pathology in the rotator cuff: a review of literature and implications for rehabilitation and surgery. *Br J Sports Med.* 2010 Sep;44(12):838–47.
 14. Neer CS 2nd. Impingement lesions. *Clin Orthop Relat Res.* 1983 Mar;(173):70–7.
 15. Bigliani LU, Ticker JB, Flatow EL, Soslowsky LJ, Mow VC. The relationship of acromial architecture to rotator cuff disease. *Clin Sports Med.* 1991 Oct;10(4):823–38.
 16. Gill TJ, McIrvin E, Kocher MS, Homa K, Mair SD, Hawkins RJ. The relative importance of acromial morphology and age with respect to rotator cuff pathology. *J Shoulder Elbow Surg.* 2002 Jul–Aug;11(4):327–30.
 17. Tamai M, Okajima S, Fushiki S, Hirasawa Y. Quantitative analysis of neural distribution in human coracoacromial ligaments. *Clin Orthop Relat Res.* 2000 Apr;(373):125–34.
 18. Edelson JG, Taitz C. Anatomy of the coraco-acromial arch. Relation to degeneration of the acromion. *J Bone Joint Surg Br.* 1992 Jul;74(4):589–94.
 19. Ogata S, Uhthoff HK. Acromial enthesopathy and rotator cuff tear. A radiologic and histologic postmortem investigation of the coracoacromial arch. *Clin Orthop Relat Res.* 1990 May;(254):39–48.
 20. Harryman DT 2nd, Hettrich CM, Smith KL, Campbell B, Sidles JA, Matsen FA 3rd. A prospective multipractice investigation of patients with full-thickness rotator cuff tears: the importance of comorbidities, practice, and other covariables on self-assessed shoulder function and health status. *J Bone Joint Surg Am.* 2003 Apr;85-A(4):690–6.
 21. Mallon WJ, Misamore G, Snead DS, Denton P. The impact of preoperative smoking habits on the results of rotator cuff repair. *J Shoulder Elbow Surg.* 2004 Mar–Apr;13(2):129–32.
 22. Ide K, Shirai Y, Ito H. Sensory nerve supply in the human subacromial bursa. *J Shoulder Elbow Surg.* 1996 Sep–Oct;5(5):371–82.
 23. Aszmann OC, Dellon AL, Birely BT, McFarland EG. Innervation of the human shoulder joint and its implications for surgery. *Clin Orthop Relat Res.* 1996 Sep;(330):202–7.
 24. Voloshin I, Gelinis J, Maloney MD, O'Keefe RJ, Bigliani LU, Blaine TA. Pro-inflammatory cytokines and metalloproteases are expressed in the subacromial bursa in patients with rotator cuff disease. *Arthroscopy.* 2005 Sep;21(9):1076.
 25. Teefey SA, Rubin DA, Middleton WD, Hildebolt CF, Leibold RA, Yamaguchi K. Detection and quantification of rotator cuff tears: comparison of ultrasonographic, magnetic resonance imaging, and arthroscopic findings in seventy-one consecutive cases. *J Bone Joint Surg Am.* 2004 Apr;86-A(4):708–16.
 26. Pedowitz RA, Yamaguchi K, Ahmad CS, et al. American Academy of Orthopaedic Surgeons. Optimizing the management of rotator cuff problems. *J Am Acad Orthop Surg.* 2011 Jun;19(6):368–79.
 27. Djurasovic M, Marra G, Arroyo JS. Revision rotator cuff repair: factors influencing results. *J Bone Joint Surg Am.* 2001 Dec;83-A(12):1849–55.
 28. Galatz LM, Griggs S, Cameron BD, Iannotti JP. Prospective longitudinal analysis of postoperative shoulder function: a ten-year follow-up study of full-thickness rotator cuff tears. *J Bone Joint Surg Am.* 2001 Jul;83-A(7):1052–6.
 29. Nho SJ, Shindle MK, Sherman SL, Freedman KB, Lyman S, MacGillivray JD. Systematic review of arthroscopic rotator cuff repair and mini-open rotator cuff repair. *J Bone Joint Surg Am.* 2007 Oct;89(Suppl 3):127–36.
 30. Trappey GJ 4th, Gartsman GM. A systematic review of the clinical outcomes of single row versus double row rotator cuff repairs. *J Shoulder Elbow Surg.* 2011 Mar;20(2 Suppl):S14–9.